

VA



U.S. Department
of Veterans Affairs

Impact of Participation in VA Evidence-Based Psychotherapy for Depression on Suicidal Ideation among Veterans


OMHSP National Evidence-Based Psychotherapy Training Program
VA Psychology Leadership Conference
May 31st, 2018



Suicidal Ideation & Mental Health

- Suicidal ideation (SI) associated with diagnoses of posttraumatic stress disorder, bipolar disorder, and major depression has been shown to predict subsequent suicide attempts (Nock et al., 2009)
- VA/DoD Clinical Practice Guidelines for Assessment and Management of Patients at Risk for Suicide (2013) recommend that “...patients receive optimal evidence-based treatment for any mental health and medical conditions that may be related to the risk of suicide”


CPG Recommended Interventions



- EBP's are consistent with recommendations at every level of suicide risk

CPG Recommended Interventions

- Veterans with a history of suicide attempts, active ideation/intent, existence of warning signs/risk factors, and limited protective factors are likely to be at high risk for suicide
- Safety Planning and suicide-focused psychotherapies (e.g., CBT for suicide prevention) are recommended for Veterans at high acute risk
- The VA EBP Training Program is currently developing advanced Suicide Safety Planning Training to be rolled out to providers who work with Veterans with elevated suicide risk
- Other VA efforts are underway to increase access to CBT for Suicide Prevention





Suicidal Ideation & EBPs for Depression

- Cognitive Behavioral Therapy for Depression (CBT-D), Acceptance and Commitment Therapy for Depression (ACT-D), and Interpersonal Psychotherapy (IPT) are identified as first-line recommended treatments for mild to moderate depression in the VA/DoD Clinical Practice Guideline (CPG) for the Management of Major Depressive Disorder (2016)
 - The evidence does not support recommending a specific evidence-based psychotherapy or pharmacotherapy over another
 - A combination of pharmacotherapy and evidence-based psychotherapy is suggested for the treatment of patients with severe, chronic, or recurrent Major Depressive Disorder



Suicidal Ideation & EBPs for Depression

- Since 2008, approximately 2,700 clinicians have been trained by VA's National Evidence-Based Psychotherapy (EBP) Training Programs in CBT-D, ACT-D and IPT
- CBT-D and ACT-D have been shown to significantly reduce SI among Veterans (Brown, Karlin, Trockel, Gordienko, Yesavage & Taylor, 2016; Walser, Garvert, Karlin, Trockel, Ryu, & Taylor, 2015)
- Additional analyses of data from these VA training programs were conducted to address the following questions:
 - Across EBPs for depression, what is the impact of treatment on Veterans' suicidal ideation?
 - Are EBPs for Depression effective in addressing moderate to severe suicidal ideation?




EBP Program Data

- Sample included **1,808 Veterans** who endorsed SI on the Beck Depression Inventory (BDI-II) at the time of the first therapy session
- Suicidal Ideation (assessed by BDI-II item 9)*
 - 91% indicated “I have thoughts of killing myself, but I would not carry them out” (score = 1)
 - 7% indicated “I would like to kill myself” (score = 2)
 - 2% indicated “I would kill myself if I had the chance” (score = 3)
- Mean BDI-II score at treatment initiation was 34.3 ($SD = 9.5$), which is in the severe range

*Depression rating scales (e.g., BDI-II), have been shown to be valid measurements of suicidal ideation (Desseilles, Perroud, Guillaume, Jausse, Genty, Malafosse, & Courtet, 2012)

Participation in EBP treatment was associated with statistically significant decreases in suicidal ideation at both the mid-point and treatment termination sessions




*All timepoints are significantly different from each other at $p < .001$.

**Last observation carried forward for missing data at mid-tx and last session.

N = 1,808 Veterans endorsing suicidal ideation who participated in IPT (*n* = 271), CBT-D (*n* = 855), or ACT (*n* = 682).


Participation in EBP treatment was associated with statistically significant decreases in suicidal ideation at mid-point and termination among the 163 Veterans who endorsed more severe suicidal ideation



*All timepoints are significantly different from each other at $p \leq .001$.

**Last observation carried forward for missing data at mid-tx and last session.

Veterans who were more severely depressed at the first session were more likely to be on the severe suicidal ideation trajectory ($\beta = .08, p < .0001$) and to experience a faster decrease in suicidal ideation over time





Summary & Limitations

- Clinical Practice Guidelines recommend first-line treatments for conditions underlying suicide, including ACT-D, CBT-D, and IPT for mild to moderate depression
- CBT-D, ACT-D, and IPT appear to decrease suicidal ideation early in treatment regardless of initial severity
 - Data is for program evaluation, not research; therefore participants were not randomized and causality cannot be demonstrated
 - Conclusions are limited by the assessment used to measure SI
 - Missing data limits conclusions
- Presence of SI should not necessarily be seen as factor for exclusion from EBP treatment



Future Directions

- Conduct additional analyses examining the impact of EBPs for other disorders and conditions associated with increased risk for suicide
 - Other VA EBPs, including Cognitive Behavioral Therapy for Insomnia (CBT-I) and Prolonged Exposure Therapy (PE), have been shown to significantly reduce suicidal ideation (SI) among Veterans
 - Cognitive Processing Therapy (CPT) has been shown to reduce SI in civilian and active duty military samples
- Future research should consider prospective follow-up after completion of EBPs to assess impacts on suicidal behavior



Contributing Authors

Kristin Powell, Ph.D.

Mandy Kumpula, Ph.D.

Kristine Day, Ph.D.

Nathan Kimbrel, Ph.D.

H. Ryan Wagner, Ph.D.

Chris Crowe, Ph.D.

Wendy Batdorf, Ph.D.

Hani Shabana, Ph.D.



U.S. Department
of Veterans Affairs

Questions/Discussion